

For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory

**MYCAMINE® 50 mg / 100 mg
Micafungin Sodium for Injection 50 mg / 100 mg**

QUALITATIVE AND QUANTITATIVE COMPOSITION

Mycamine 50 mg powder for solution for Injection:

Each vial contains:
Micafungin Sodium equivalent to 50 mg Micafungin.

After reconstitution each ml contains Micafungin Sodium equivalent to 10 mg Micafungin.

Mycamine 100 mg powder for solution for Injection:

Each vial contains:
Micafungin Sodium equivalent to 100 mg Micafungin.

After reconstitution each ml contains Micafungin Sodium equivalent to 20 mg Micafungin.

PHARMACEUTICAL FORM

Powder for solution for infusion.

White compact powder.

CLINICAL PARTICULARS

Therapeutic Indications

MYCAMINE is indicated for:

- **Treatment of Patients with Candidemia, Acute Disseminated Candidiasis, *Candida* Peritonitis and Abscesses.**
MYCAMINE has not been adequately studied in patients with endocarditis, osteomyelitis and meningitis due to *Candida* infections.
- **Treatment of Patients with Esophageal Candidiasis.**
- **Prophylaxis of *Candida* and *Aspergillus* Infections in Patients Undergoing Hematopoietic Stem Cell Transplantation.**
- **Treatment of Patients with Fungemia, Respiratory mycosis, Gastrointestinal mycosis Caused by *Aspergillus* sp..**

Note: The efficacy of *MYCAMINE* against infections caused by fungi other than *Candida* sp. and *Aspergillus* sp. has not been established.

Posology and Method of Administration

Do not mix or co-infuse *MYCAMINE* with other medications. *MYCAMINE* has been shown to

precipitate when mixed directly with a number of other commonly used medications.

Table 1:MYCAMINE Dosage

Indication	Recommended Reconstituted Dose Once Daily
Treatment of Candidemia, Acute Disseminated Candidiasis, <i>Candida</i> Peritonitis and Abscesses ¹	100 mg
Treatment of Esophageal Candidiasis ²	150 mg
Prophylaxis of <i>Candida</i> and <i>Aspergillus</i> Infections in Hematopoietic Stem Cell Transplant (HSCT) recipients ³	50 mg
Treatment of <i>Aspergillus</i> Infections (Fungemia, respiratory mycosis, gastrointestinal mycosis) ^{4,5}	100 – 150 mg

¹ In patients treated successfully for candidemia and other *Candida* infections, the mean duration of treatment was 15 days (range 10-47 days).

² In patients treated successfully for esophageal candidiasis, the mean duration of treatment was 15 days (range 10-30 days).

³ In hematopoietic stem cell transplant (HSCT) recipients who experienced success of prophylactic therapy, the mean duration of prophylaxis was 19 days (range 6-51 days).

⁴ For severe or refractory aspergillosis, the dosage can be increased up to 300 mg /day according to the patient's condition.

⁵ Combination therapy of Mycamine with other antifungals has also been tried in clinical trials.

A loading dose is not required. Typically, 85% of the steady-state concentration is achieved after three daily *MYCAMINE* doses.

No dosing adjustments are required based on race, gender, or in patients with severe renal impairment or in patients with mild, moderate or severe hepatic impairment.

No dose adjustment for *MYCAMINE* is required with concomitant use of mycophenolate mofetil, cyclosporine, tacrolimus, prednisolone, sirolimus, nifedipine, fluconazole, voriconazole, itraconazole, amphotericin B, ritonavir, or rifampin.

Directions for Reconstitution and Dilution

Please read this entire section carefully before beginning reconstitution.

The diluent to be used for reconstitution and dilution is 0.9% Sodium Chloride solution for infusion. Alternatively, 5% Dextrose solution for infusion, may be used for reconstitution and dilution of *MYCAMINE*. Solutions for infusion are prepared as follows:

Reconstitution

Aseptically add 5 ml of 0.9% Sodium Chloride solution for infusion or 5 ml of 5% Dextrose solution for infusion to each 50 mg vial to yield a preparation containing approximately 10 mg micafungin/ml.

Aseptically add 5 ml of 0.9% Sodium Chloride solution for infusion or 5 ml of 5% Dextrose solution for infusion to each 100 mg vial to yield a preparation containing approximately 20 mg micafungin/ml.

As with all parenteral drug products, reconstituted *MYCAMINE* should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use material if there is any evidence of precipitation or foreign matter. Aseptic technique must be strictly observed in all handling since no preservative or bacteriostatic agent is present in *MYCAMINE* or in the materials specified for reconstitution and dilution.

Dissolution

To minimize excessive foaming, GENTLY dissolve the *MYCAMINE* powder by swirling the vial. DO NOT VIGOROUSLY SHAKE THE VIAL. Visually inspect the vial for particulate matter.

Dilution

The diluted solution should be protected from light. It is not necessary to cover the infusion drip chamber or the tubing.

- For treatment of candidemia, acute disseminated candidiasis, *Candida* peritonitis and abscesses :
add 100 mg of reconstituted *MYCAMINE* into 100 ml of 0.9% Sodium Chloride solution for infusion or 100 ml of 5% Dextrose solution for infusion.
- For treatment of esophageal candidiasis:
add 150 mg of reconstituted *MYCAMINE* into 100 ml of 0.9% Sodium Chloride solution for infusion or 100 ml of 5% Dextrose solution for infusion.
- Treatment of *Aspergillus* infections:
Add 100 to 150 mg of reconstituted Mycamine into 100 ml of 0.9% Sodium Chloride solution or 100 ml of 5% Dextrose solution for infusion
- For prophylaxis of *Candida* and *Aspergillus* infections:
add 50 mg of reconstituted *MYCAMINE* into 100 ml of 0.9% Sodium Chloride solution for infusion or 100 ml of 5% Dextrose solution for infusion.

MYCAMINE is preservative-free. Discard partially used vials.

Infusion Volume and Duration

MYCAMINE should be administered by intravenous infusion only. Infuse over one hour. More rapid infusions may result in more frequent histamine mediated reactions.

An existing intravenous line should be flushed with 0.9% Sodium Chloride solution for infusion prior to infusion of *MYCAMINE*.

Use in Specific Populations

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

A total of 418 subjects in clinical studies of *MYCAMINE* were 65 years of age and older, and 124 subjects were 75 years of age and older. No overall differences in safety and effectiveness were observed between these subjects and younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

The exposure and disposition of a 50 mg *MYCAMINE* dose administered as a single 1-hour infusion to 10 healthy subjects aged 66-78 years were not significantly different from those in 10 healthy subjects aged 20-24 years. No dose adjustment is necessary for the elderly.

Use in Patients with Renal Impairment

MYCAMINE does not require dose adjustment in patients with renal impairment. Supplementary dosing should not be required following hemodialysis

Use in Patients with Hepatic Impairment

Dose adjustment of *MYCAMINE* is not required in patients with mild, moderate or severe hepatic impairment.

Race and Gender

No dose adjustment of *MYCAMINE* is required based on gender or race. After 14 daily doses of 150 mg to healthy subjects, micafungin AUC in women was greater by approximately 23% compared with men, due to smaller body weight. No notable differences among white, black, and Hispanic subjects were seen. The micafungin AUC was greater by 19% in Japanese subjects compared to blacks, due to smaller body weight.

Drug Abuse and Dependence

There has been no evidence of either psychological or physical dependence, or withdrawal or rebound effects with *MYCAMINE*.

Contraindications

MYCAMINE is contraindicated in persons with known hypersensitivity to micafungin, any component of *MYCAMINE*, or other echinocandins.

Special Warnings and Special Precautions for Use

Hypersensitivity Reactions

Isolated cases of serious hypersensitivity (anaphylaxis and anaphylactoid) reactions (including shock) have been reported in patients receiving *MYCAMINE*. If these reactions occur, *MYCAMINE* infusion should be discontinued and appropriate treatment administered.

Hematological Effects

Acute intravascular hemolysis and hemoglobinuria was seen in a healthy volunteer during infusion of *MYCAMINE* (200 mg) and oral prednisolone (20 mg). This reaction was transient, and the subject did not develop significant anemia. Isolated cases of significant hemolysis and hemolytic anemia have also been reported in patients treated with *MYCAMINE*. Patients who develop clinical or laboratory evidence of hemolysis or hemolytic anemia during *MYCAMINE* therapy should be monitored closely for evidence of worsening of these conditions and evaluated for the risk/benefit of continuing *MYCAMINE* therapy.

Hepatic Effects

Laboratory abnormalities in liver function tests have been seen in healthy volunteers and patients treated with *MYCAMINE*. In some patients with serious underlying conditions who were receiving *MYCAMINE* along with multiple concomitant medications, clinical hepatic abnormalities have occurred, and isolated cases of significant hepatic impairment, hepatitis, and hepatic failure have been reported. Patients who develop abnormal liver function tests during *MYCAMINE* therapy should be monitored for evidence of worsening hepatic function and evaluated for the risk/benefit of continuing *MYCAMINE* therapy.

Renal Effects

Elevations in BUN and creatinine, and isolated cases of significant renal impairment or acute renal failure have been reported in patients who received *MYCAMINE*. In fluconazole-controlled trials, the incidence of drug-related renal adverse events was 0.4% for *MYCAMINE* treated patients and 0.5% for fluconazole treated patients. Patients who develop abnormal renal function tests during *MYCAMINE* therapy should be monitored for evidence of worsening renal function.

Interaction with Other Medicaments and Other Forms of Interaction

A total of 14 clinical drug-drug interaction studies were conducted in healthy volunteers to evaluate the potential for interaction between *MYCAMINE* and amphotericin B, mycophenolate mofetil, cyclosporine, tacrolimus, prednisolone, sirolimus, nifedipine, fluconazole, itraconazole, voriconazole, ritonavir, and rifampin. In these studies, no interaction that altered the pharmacokinetics of micafungin was observed.

There was no effect of a single dose or multiple doses of *MYCAMINE* on mycophenolate mofetil, cyclosporine, tacrolimus, prednisolone, fluconazole, and voriconazole pharmacokinetics.

Sirolimus AUC was increased by 21% with no effect on C_{max} in the presence of steady-state *MYCAMINE* compared with sirolimus alone. Nifedipine AUC and C_{max} were increased by 18% and 42%, respectively, in the presence of steady-state *MYCAMINE* compared with nifedipine alone. Itraconazole AUC and C_{max} were increased by 22% and 11%, respectively.

Patients receiving sirolimus, nifedipine or itraconazole in combination with *MYCAMINE* should be monitored for sirolimus, nifedipine or itraconazole toxicity and the sirolimus, nifedipine or itraconazole dosage should be reduced if necessary.

Micafungin is neither a substrate nor an inhibitor of P-glycoprotein and, therefore, would not be expected to alter P-glycoprotein-mediated drug transport activity.

Pregnancy and Lactation

Pregnancy

There are no adequate and well-controlled studies of micafungin in pregnant women. Animal reproduction studies in rabbits showed visceral abnormalities and increased abortion at 4 times the recommended human dose. However, animal studies are not always predictive of human response.

Micafungin should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

When pregnant rabbits were given 4 times the recommended human dose, there were increased abortion and visceral abnormalities including abnormal lobation of the lung, levocardia, retrocaval ureter, anomalous right subclavian artery, and dilatation of the ureter.

Lactation

It is not known whether micafungin is excreted in human milk. Caution should be exercised when *MYCAMINE* is administered to a nursing woman.

Effects on Ability to Drive and Use Machines

No studies on the effects on the ability to drive and use machines have been performed. However, adverse reactions may occur, which may influence the ability to drive and use machines (see *Undesirable Effects*).

Undesirable Effects

General

Possible histamine-mediated symptoms have been reported with *MYCAMINE*, including rash, pruritus, facial swelling, and vasodilatation.

Injection site reactions, including phlebitis and thrombophlebitis have been reported, at *MYCAMINE* doses of 50-150 mg/day. These reactions tended to occur more often in patients receiving *MYCAMINE* via peripheral intravenous administration.

Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in clinical trials of *MYCAMINE* cannot be directly compared to rates in clinical trials of another drug and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does provide a basis for identifying adverse events that appear to be related to drug use and for approximating rates.

Candidemia and Other Candida Infections

In a randomized, double-blind study for treatment of candidemia and other *Candida* infections, treatment emergent adverse events occurred in 183/200 (91.5%), 187/202 (92.6%) and 171/193 (88.6%) patients in the *MYCAMINE* 100 mg/day, *MYCAMINE* 150mg/day, and caspofungin (70/50mg/day) treatment groups, respectively.

Treatment emergent adverse events occurring in $\geq 5\%$ of the patients in any treatment study groups are shown in Table 2.

Table 2: *Treatment Emergent Adverse Events in Patients with Candidemia and Other *Candida* Infections

Adverse Event ⁽¹⁾ MedDRA v 5.0 System Organ Class Preferred Term	Micafungin 100 mg (n = 200)	Micafungin 150 mg (n = 202)	Caspofungin ⁽²⁾ (n = 193)
All Systems, Any Adverse Event	183 (91.5)	187 (92.6)	171 (88.6)
Gastrointestinal Disorders	81 (40.5)	89 (44.1)	76 (39.4)
Diarrhea NOS	15 (7.5)	26 (12.9)	14 (7.3)
Nausea	19 (9.5)	15 (7.4)	20 (10.4)
Vomiting NOS	18 (9)	15 (7.4)	16 (8.3)
Abdominal Pain NOS	5 (2.5)	4 (2)	10 (5.2)
Metabolism and Nutrition Disorders	77 (38.5)	83 (41.1)	73 (37.8)
Hypokalemia	28 (14)	34 (16.8)	28 (14.5)
Hypomagnesaemia	11 (5.5)	17 (8.4)	14 (7.3)
Hypoglycemia NOS	12 (6)	14 (6.9)	9 (4.7)
Hypernatremia	8 (4)	13 (6.4)	8 (4.1)
Hyperkalemia	10 (5)	8 (4)	5 (2.6)
Infections and Infestations	67 (33.5)	81 (40.1)	59 (30.6)
Bacteremia	10 (5)	18 (8.9)	11 (5.7)
Septic Shock	15 (7.5)	9 (4.5)	9 (4.7)
Sepsis NOS	11 (5.5)	10 (5)	11 (5.7)
Pneumonia NOS	3 (1.5)	11 (5.4)	4 (2.1)
General Disorders / Administration Site Conditions	59 (29.5)	56 (27.7)	51 (26.4)
Pyrexia	14 (7)	22 (10.9)	15 (7.8)
Edema Peripheral	11 (5.5)	12 (5.9)	14 (7.3)
Vascular Disorders	43 (21.5)	47 (23.3)	36 (18.7)
Hypotension NOS	20 (10)	12 (5.9)	15 (7.8)
Hypertension NOS	6 (3)	10 (5)	12 (6.2)
Investigations	36 (18)	49 (24.3)	37 (19.2)
Blood Alkaline Phosphatase NOS Increased	11 (5.5)	16 (7.9)	8 (4.1)
Blood /Lymphatic System Disorders	38 (19)	45 (22.3)	37 (19.2)

Thrombocytopenia	8 (4)	8 (4)	11 (5.7)
Anemia NOS	5 (2.5)	6 (3)	13 (6.7)
Anemia NOS Aggravated	4 (2)	10 (5)	5 (2.6)
Cardiac Disorders	35 (17.5)	48 (23.8)	36 (18.7)
Tachycardia NOS	6 (3)	7 (3.5)	13 (6.7)
Bradycardia NOS	5 (2.5)	10 (5)	8 (4.1)
Atrial Fibrillation	5 (2.5)	10 (5)	0
Nervous System Disorders	21 (10.5)	42 (20.8)	32 (16.6)
Headache NOS	4 (2)	10 (5)	11 (5.7)
Skin/Subcutaneous Tissue Disorders	26 (13)	34 (16.8)	33 (17.1)
Decubitus Ulcer	9 (4.5)	12 (5.9)	9 (4.7)
Psychiatric Disorders	31 (15.5)	27 (13.4)	33 (17.1)
Insomnia	11 (5.5)	8 (4)	16 (8.3)

Patient base: all randomized patients who received at least 1 dose of trial drug

Common: $\geq 5\%$ in any treatment arm.

*During IV treatment + 3 day

⁽¹⁾ Within a system organ class patients may experience more than 1 adverse event.

⁽²⁾ 70 mg loading dose on day 1 followed by 50 mg/day thereafter (casposfungin)

In a second, supportive, randomized, double-blind study for treatment of candidemia and other *Candida* infections, treatment emergent adverse events occurred in 245/264 (92.8%) and 250/265 (94.3%) patients in the *MYCAMINE* (100 mg/day) and AmBisome (3 mg/kg/day) treatment groups, respectively. The most common treatment emergent adverse events occurring in $\geq 5\%$ of the *MYCAMINE*-treated patients at least 16 years of age were: pyrexia (15.2% vs. 17%); hypokalemia (16.7% vs. 20.8%); nausea (9.5% vs. 8.3%); diarrhea (10.6% vs. 11.3%) and vomiting (12.9% vs. 9.4%), in the *MYCAMINE* and AmBisome treatment groups, respectively. Other important treatment emergent adverse events that occurred at $< 5\%$ frequency were abnormal liver function tests (4.2% vs. 3%); increased aspartate aminotransferase (2.7% vs. 1.9%), and increased blood alkaline phosphatase (3% vs. 2.3%), in the *MYCAMINE* and AmBisome treatment groups, respectively.

Esophageal Candidiasis

In a randomized, double-blind study for treatment of esophageal candidiasis, a total of 202/260 (77.7%) patients who received *MYCAMINE* 150 mg/day and 186/258 (72.1%) patients who received intravenous fluconazole 200 mg/day experienced an adverse event. Treatment emergent adverse events resulting in discontinuation were reported in 17 (6.5%) *MYCAMINE* treated patients; and in 12 (4.7%) fluconazole treated patients. Treatment emergent adverse events occurring in $\geq 5\%$ of the patients in either treatment group are shown in Table 3.

Table 3: *Treatment Emergent Adverse Events in Patients with Esophageal Candidiasis

Adverse Events ⁽¹⁾ (MedDRA System Organ Class and Preferred Term)	<i>MYCAMINE</i> 150 mg/day n (%)	Fluconazole 200 mg/day n (%)
Number of Patients	260	258
All Systems, Any Adverse Event	202 (77.7)	186 (72.1)
Gastrointestinal Disorders	84 (32.3)	93 (36)
Diarrhea NOS	27 (10.4)	29 (11.2)
Nausea	20 (7.7)	23 (8.9)
Vomiting NOS	17 (6.5)	17 (6.6)
Abdominal Pain NOS	10 (3.8)	15 (5.8)
General Disorders / Administration Site Conditions	52 (20)	45 (17.4)
Pyrexia	34 (13.1)	21 (8.1)
Nervous System Disorders	42 (16.2)	40 (15.5)
Headache NOS	22 (8.5)	20 (7.8)
Blood /Lymphatic System Disorders	38 (14.6)	43 (16.7)
Anemia NOS	8 (3.1)	16 (6.2)
Vascular Disorders	54 (20.8)	21 (8.1)
Phlebitis NOS	49 (18.8)	13 (5)
Skin and Subcutaneous Tissue Disorders	36 (13.8)	26 (10.1)
Rash NOS	14 (5.4)	6 (2.3)
Psychiatric Disorders	20 (7.7)	21 (8.1)
Insomnia	9 (3.5)	13 (5)

Patient base: all randomized patients who received at least 1 dose of trial drug.

Common: $\geq 5\%$ in either treatment arm.

* During treatment + 3 days.

⁽¹⁾ Within a system organ class patients may experience more than 1 adverse event.

Prophylaxis of Candida and Aspergillus Infections in Hematopoietic Stem Cell Transplant Recipients

A double-blind study was conducted in a total of 882 patients scheduled to undergo an autologous or allogeneic hematopoietic stem cell transplant. The median duration of treatment was 18 days (range 1 to 51 days) in both treatment arms.

All patients who received *MYCAMINE* (425) and all patients who received fluconazole (457) experienced at least one adverse event during the study. Treatment emergent adverse events resulting in *MYCAMINE* discontinuation were reported in 18 (4.2%) patients; while those resulting in fluconazole discontinuation were reported in 33 (7.2%). Treatment emergent adverse events occurring in $\geq 15\%$ of the patients in either treatment group are shown in Table 4.

Table 4: * Treatment Emergent Adverse Events During Prophylaxis of *Candida* and *Aspergillus* Infection in Hematopoietic Stem Cell Transplant Recipients

Adverse Events ⁽¹⁾ (MedDRA System Organ Class and Preferred Term)	<i>MYCAMINE</i> 50 mg/day n (%)	Fluconazole 400 mg/day n (%)
Number of Patients	425	457
All Systems, Any Adverse Events	425 (100)	457 (100)
Gastrointestinal Disorders	421 (99.1)	449 (98.2)
Diarrhea NOS	302 (71.1)	348 (76.1)
Nausea	296 (69.6)	309 (67.6)
Vomiting NOS	281 (66.1)	307 (67.2)
Constipation	129 (30.4)	143 (31.3)
Dyspepsia	104 (24.5)	122 (26.7)
Abdominal Pain NOS	115 (27.1)	107 (23.4)
General Disorders / Administration Site Conditions	410 (96.5)	440 (96.3)
Mucosal Inflammation NOS	322 (75.8)	360 (78.8)
Pyrexia	191 (44.9)	218 (47.7)
Fatigue	126 (29.6)	145 (31.7)
Rigors	112 (26.4)	118 (25.8)
Edema Peripheral	88 (20.7)	100 (21.9)
Blood and Lymphatic System Disorders	408 (96)	429 (93.9)
Neutropenia	320 (75.3)	327 (71.6)
Thrombocytopenia	307 (72.2)	304 (66.5)
Anemia NOS	151 (35.5)	173 (37.9)
Febrile Neutropenia	155 (36.5)	166 (36.3)
Metabolism and Nutrition Disorders	385 (90.6)	428 (93.7)
Hypomagnesaemia	214 (50.4)	256 (56)
Hypokalemia	209 (49.2)	232 (50.8)
Anorexia	116 (27.3)	121 (26.5)
Appetite Decreased NOS	87 (20.5)	93 (20.4)
Fluid Overload	74 (17.4)	96 (21)
Hyperglycemia NOS	68 (16)	92 (20.1)
Hypocalcemia	72 (16.9)	82 (17.9)
Fluid Retention	69 (16.2)	66 (14.4)
Respiratory, Thoracic and Mediastinal Disorders	291 (68.5)	336 (73.5)
Cough	98 (23.1)	112 (24.5)
Epistaxis	49 (11.5)	84 (18.4)
Dyspnea NOS	54 (12.7)	64 (14)
Skin and Subcutaneous Tissue Disorders	290 (68.2)	316 (69.1)
Rash NOS	110 (25.9)	102 (22.3)
Pruritus NOS	75 (17.6)	87 (19)
Erythema	48 (11.3)	71 (15.5)
Nervous System Disorders	261 (61.4)	268 (58.6)
Headache NOS	179 (42.1)	165 (36.1)

Dizziness	55 (12.9)	83 (18.2)
Psychiatric Disorders	257 (60.5)	249 (54.5)
Insomnia	152 (35.8)	146 (31.9)
Anxiety	95 (22.4)	92 (20.1)
Vascular Disorders	224 (52.7)	267 (58.4)
Hypertension NOS	91 (21.4)	113 (24.7)
Hypotension NOS	79 (18.6)	89 (19.5)
Flushing	47 (11.1)	70 (15.3)
Infections and Infestations	178 (41.9)	208 (45.5)
Bacteremia	66 (15.5)	86 (18.8)
Cardiac Disorders	147 (34.6)	162 (35.4)
Tachycardia NOS	105 (24.7)	102 (22.3)

Patient base: all randomized patients who received at least 1 dose of trial drug.

Common: $\geq 5\%$ in either treatment arm.

* During treatment + 3 days.

⁽¹⁾ Within a system organ class patients may experience more than 1 adverse event.

Overall MYCAMINE Safety Experience in Clinical Trials

The overall safety of *MYCAMINE* was assessed in 3083 patients and 501 volunteers in 41 clinical studies, including the invasive candidiasis, esophageal candidiasis, aspergillus infections and prophylaxis studies, who received single or multiple doses of *MYCAMINE*, ranging from 12.5 mg to ≥ 150 mg/day. Treatment emergent adverse events which occurred in $\geq 5\%$ of all patients who received *MYCAMINE* in these trials are shown in Table 5.

Overall, 2810 of 3083 (91.1%) patients who received *MYCAMINE* experienced an adverse event.

Clinically significant adverse events regardless of causality or incidence which occurred in these trials are listed below:

- *Blood and lymphatic system disorders*: coagulopathy, febrile neutropenia, hemolysis, hemolytic anemia, pancytopenia, thrombotic thrombocytopenic purpura
- *Cardiac disorders*: arrhythmia, atrial fibrillation, cardiac arrest, cyanosis, hypotension, myocardial infarction, tachycardia
- *Gastrointestinal disorders*: abdominal pain upper, dyspepsia
- *General disorders and administration site conditions*: injection site thrombosis
- *Hepatobiliary disorders*: hepatocellular damage, hepatomegaly, jaundice, hepatic failure
- *Infections and infestations*: infection, pneumonia, sepsis
- *Metabolism and nutrition disorders*: acidosis, anorexia, hyponatremia
- *Musculoskeletal, connective tissue and bone disorders*: arthralgia
- *Nervous system disorders*: convulsions, encephalopathy, intracranial hemorrhage
- *Psychiatric disorders*: delirium
- *Renal and urinary disorders*: anuria, hemoglobinuria, oliguria, renal failure acute, renal tubular necrosis

- *Respiratory, thoracic and mediastinal disorders*: apnea, dyspnea, hypoxia, pulmonary embolism
- *Skin and subcutaneous tissue disorders*: erythema multiforme, skin necrosis, urticaria
- *Vascular disorders*: deep venous thrombosis, hypertension

Table 5: *Treatment Emergent Adverse Events in Patients Who Received MYCAMINE in Clinical Trials

Adverse Events ⁽¹⁾ (MedDRA System Organ Class and Preferred Term)	MYCAMINE n (%)
Number of Patients	3083
All Systems, Any Adverse Event	2810 (91.1)
Gastrointestinal Disorders	1764 (57.2)
Diarrhea NOS	718 (23.3)
Nausea	679 (22)
Vomiting NOS	669 (21.7)
Constipation	341 (11.1)
Abdominal Pain	300 (9.7)
Dyspepsia	176 (5.7)
General Disorders / Administration Site Conditions	1407 (45.6)
Pyrexia	618 (20)
Mucosal Inflammation NOS	438 (14.2)
Rigors	281 (9.1)
Edema Peripheral	209 (6.8)
Fatigue	198 (6.4)
Metabolism and Nutrition Disorders	1316 (42.7)
Hypokalemia	556 (18)
Hypomagnesemia	409 (13.3)
Hypocalcemia	201 (6.5)
Anorexia	190 (6.2)
Hyperglycemia NOS	173 (5.6)
Fluid Overload	155 (5)
Infections and Infestations	1227 (39.8)
Bacteremia	185 (6)
Sepsis NOS	156 (5.1)
Respiratory, Thoracic and Mediastinal Disorders	1108 (35.9)
Cough	251 (8.1)
Dyspnea NOS	182 (5.9)
Epistaxis	172 (5.6)
Blood and Lymphatic System Disorders	1047 (34)
Thrombocytopenia	474 (15.4)
Neutropenia	436 (14.1)
Anemia NOS	302 (9.8)
Febrile Neutropenia	187 (6.1)
Investigations	989 (32.1)
Aspartate Aminotransferase Increased	172 (5.6)

Adverse Events ⁽¹⁾ (MedDRA System Organ Class and Preferred Term)	MYCAMINE n (%)
Blood Alkaline Phosphatase NOS Increased	168 (5.4)
Alanine Aminotransferase Increased	165 (5.4)
Skin and Subcutaneous Tissue Disorders	940 (30.5)
Rash NOS	269 (8.7)
Pruritus NOS	187 (6.1)
Nervous System Disorders	889 (28.8)
Headache NOS	489 (15.9)
Psychiatric Disorders	727 (23.6)
Insomnia	303 (9.8)
Anxiety	198 (6.4)
Vascular Disorders	867 (28.1)
Hypotension NOS	279 (9.1)
Hypertension NOS	214 (6.9)
Phlebitis NOS	172 (5.6)
Musculoskeletal and Connective Tissue Disorders	579 (18.8)
Back Pain	166 (5.4)
Cardiac Disorders	563 (18.3)
Tachycardia NOS	231 (7.5)

Patient base: all randomized patients who received at least 1 dose of trial drug.

Common: Incidence of adverse event $\geq 5\%$.

* During treatment + 3 days.

⁽¹⁾ Within a system organ class, patients may experience more than 1 adverse event.

Postmarketing Adverse Reactions

The following adverse reactions have been identified during the post-approval use of micafungin sodium for injection. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency. A causal relationship to micafungin sodium for injection could not be excluded for these adverse reactions, which included:

- *Hepatobiliary disorders*: hyperbilirubinemia, hepatic function abnormal, hepatic disorder, hepatocellular damage
- *Renal and urinary disorders*: acute renal failure and renal impairment
- *Blood and lymphatic system disorders*: white blood cell count decreased, hemolytic anemia, disseminated intravascular coagulation.
- *Skin and subcutaneous tissue disorders*: Stevens-Johnson syndrome, toxic epidermal necrolysis.
- *Vascular disorders*: shock

Overdose

MYCAMINE is highly protein bound and, therefore, is not dialyzable. No cases of MYCAMINE overdosage have been reported. Repeated daily doses up to 8 mg/kg (maximum total dose of 896 mg) in adult patients have been administered in clinical trials with no reported dose-limiting toxicity. The minimum lethal dose of MYCAMINE is 125 mg/kg in

rats, equivalent to 8.1 times the recommended human clinical dose for esophageal candidiasis based on body surface area comparisons.

DESCRIPTION


MYCAMINE is a sterile, lyophilized product for intravenous (IV) infusion that contains micafungin sodium. Micafungin sodium is a semisynthetic lipopeptide (echinocandin) synthesized by a chemical modification of a fermentation product of *Coleophoma empetri* F-11899. Micafungin inhibits the synthesis of 1, 3-β-D-glucan, an integral component of the fungal cell wall.

Each single-use vial contains 50 mg micafungin sodium, 200 mg lactose monohydrate, with citric acid and/or sodium hydroxide (used for pH adjustment). *MYCAMINE* must be diluted with 0.9% Sodium Chloride solution for infusion or 5% Dextrose solution for infusion (refer to *Posology and Method of Administration*). Following reconstitution with 0.9% Sodium Chloride solution for infusion the resulting pH of the solution is between 5-7.

Micafungin sodium is chemically designated as:

Pneumocandin A0, 1-[(4R,5R)-4,5-dihydroxy-N2-[4-[5-[4-(pentyloxy)phenyl]-3-isoxazolyl]benzoyl]-L-ornithine]-4-[(4S)-4-hydroxy-4-[4-hydroxy-3-(sulfooxy)phenyl]-L-threonine]-, monosodium salt.

The chemical structure of micafungin sodium is:



The empirical/molecular formula is $C_{56}H_{70}N_9NaO_{23}S$ and the formula weight is 1292.26.

Micafungin sodium is a light-sensitive, hygroscopic white powder that is freely soluble in water, isotonic sodium chloride solution, N,N-dimethylformamide and dimethylsulfoxide, slightly soluble in methyl alcohol, and practically insoluble in acetonitrile, ethyl alcohol (95%), acetone, diethyl ether and n-hexane.

CLINICAL PHARMACOLOGY

Mechanism Of Action

Micafungin is a member of the echinocandin class of antifungal agents.

Pharmacokinetics

The pharmacokinetics of micafungin were determined in healthy subjects, hematopoietic stem cell transplant recipients, and patients with esophageal candidiasis up to a maximum daily dose of 8 mg/kg body weight.

The relationship of area under the concentration-time curve (AUC) to micafungin dose was linear over the daily dose range of 50 mg to 150 mg and 3 mg/kg to 8 mg/kg body weight.

Steady-state pharmacokinetic parameters in relevant patient populations after repeated daily administration are presented in the table below.

Table 6: Pharmacokinetic Parameters of Micafungin in Adult Patients

Population	n	Dose (mg)	Pharmacokinetic Parameters (Mean ± Standard Deviation)			
			C _{max} (mcg/mL)	AUC ₀₋₂₄ ¹ (mcg·h/mL)	t _{1/2} (h)	Cl (mL/min/kg)
Patients with IC	20	100	5.7±2.2	83±51	14.5±7.0	0.359 ±0.179
[Day 1] [Steady State]	20	100	10.1±4.4	97±29	13.4±2.0	0.298 ±0.115
HIV – Positive Patients with EC [Day 1] [Day 14 or 21]	20	50	4.1±1.4	36±9	14.9±4.3	0.321 ±0.098
	20	100	8.0±2.4	108±31	13.8±3.0	0.327 ±0.093
	14	150	11.6±3.1	151±45	14.1±2.6	0.340 ±0.092
	20	50	5.1±1.0	54±13	15.6±2.8	0.300±0.063
	20	100	10.1±2.6	115±25	16.9±4.4	0.301±0.086
	14	150	16.4±6.5	167±40	15.2±2.2	0.297±0.081
HSCT Recipients [Day 7]		per kg				
	8	3	21.1±2.84	234±34	14.0±1.4	0.214±0.031
	10	4	29.2±6.2	339±72	14.2±3.2	0.204±0.036
	8	6	38.4±6.9	479±157	14.9±2.6	0.224±0.064
	8	8	60.8±26.9	663±212	17.2±2.3	0.223±0.081

IC = candidemia or other *Candida* Infections; HIV = human immunodeficiency virus; EC = esophageal candidiasis; HSCT = hematopoietic stem cell transplant

¹ = AUC_{0-infinity} is presented for day 1; AUC₀₋₂₄ is presented for steady state.

Patients with Renal Impairment

MYCAMINE does not require dose adjustment in patients with renal impairment. A single 1-hour infusion of 100 mg MYCAMINE was administered to 9 subjects with severe renal impairment (creatinine clearance <30 mL/min) and to 9 age-, gender-, and weight-matched subjects with normal renal function (creatinine clearance >80 mL/min). The maximum concentration (C_{max}) and AUC were not significantly altered by severe renal impairment.

Since micafungin is highly protein bound, it is not dialyzable. Supplementary dosing should not be required following hemodialysis.

Patients with Hepatic Impairment

- A single 1-hour infusion of 100 mg *MYCAMINE* was administered to 8 subjects with moderate hepatic impairment (Child-Pugh score 7-9) and 8 age-, gender-, and weight-matched subjects with normal hepatic function. The C_{max} and AUC values of micafungin were lower by approximately 22% in subjects with moderate hepatic impairment compared to normal subjects. This difference in micafungin exposure does not require dose adjustment of *MYCAMINE* in patients with moderate hepatic impairment.
-
- A single 1-hour infusion of 100 mg *MYCAMINE* was administered to 8 subjects with severe hepatic impairment (Child-Pugh score 10-12) and 8 age-, gender-, ethnic- and weight matched subjects with normal hepatic function. The mean C_{max} and AUC values of micafungin were lower by approximately 30% in subjects with severe hepatic impairment compared to normal subjects. The mean C_{max} and AUC values of M-5 metabolite were approximately 2.3-fold higher in subjects with severe hepatic impairment compared to normal subjects; however, this exposure (parent and metabolite) was comparable to that in patients with systemic *Candida* infection. Therefore, no micafungin dose adjustment is necessary in patients with severe hepatic impairment.

Distribution

The mean \pm standard deviation volume of distribution of micafungin at terminal phase was 0.39 ± 0.11 L/kg body weight when determined in adult patients with esophageal candidiasis at the dose range of 50 mg to 150 mg.

Micafungin is highly (>99%) protein bound *in vitro*, independent of plasma concentrations over the range of 10 to 100 mcg/mL. The primary binding protein is albumin; however, micafungin, at therapeutically relevant concentrations, does not competitively displace bilirubin binding to albumin. Micafungin also binds to a lesser extent to α 1-acid-glycoprotein.

Metabolism

Micafungin is metabolized to M-1 (catechol form) by arylsulfatase, with further metabolism to M-2 (methoxy form) by catechol-O-methyltransferase. M-5 is formed by hydroxylation at the side chain (ω -1 position) of micafungin catalyzed by cytochrome P450 (CYP) isozymes. Even though micafungin is a substrate for and a weak inhibitor of CYP3A *in vitro*, hydroxylation by CYP3A is not a major pathway for micafungin metabolism *in vivo*. Micafungin is neither a P-glycoprotein substrate nor inhibitor *in vitro*.

In four healthy volunteer studies, the ratio of metabolite to parent exposure (AUC) at a dose of 150 mg/day was 6% for M-1, 1% for M-2, and 6% for M-5. In patients with esophageal candidiasis, the ratio of metabolite to parent exposure (AUC) at a dose of 150 mg/day was 11% for M-1, 2% for M-2, and 12% for M-5.

Excretion

The excretion of radioactivity following a single intravenous dose of ¹⁴C-micafungin sodium for injection (25 mg) was evaluated in healthy volunteers. At 28 days after administration, mean urinary and fecal recovery of total radioactivity accounted for 82.5% (76.4 to 87.9%) of the administered dose. Fecal excretion is the major route of elimination (total radioactivity at 28 days was 71% of the administered dose).

Microbiology

Mechanism of Action

Micafungin inhibits the synthesis of 1,3-β-D-glucan, an essential component of fungal cell walls, which is not present in mammalian cells.

Activity In Vitro

Micafungin exhibited in-vitro activity against *C. albicans*, *C. glabrata*, *C. guilliermondii*, *C. krusei*, *C. parapsilosis*, *C. tropicalis* *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus niger*, *Aspergillus terreus* and *Aspergillus nidulans*.

Standardized susceptibility testing methods for 1,3-β-D-glucan synthesis inhibitors have recently been proposed by the CLSI, however, the correlation between the results of susceptibility studies and clinical outcome has not been established.

Activity In Vivo

Micafungin sodium has shown activity in both mucosal and disseminated murine models of candidiasis. Micafungin sodium, administered to immunosuppressed mice in models of disseminated candidiasis prolonged survival and/or decreased the mycological burden.

Micafungin has been studied in murine and rabbit models of invasive aspergillosis. In these studies, micafungin was found to improve the outcome and prolong survival.

Drug Resistance

Mutants of *Candida* with reduced susceptibility to micafungin have been identified in some patients during treatment suggesting a potential for development of drug resistance. The incidence of drug resistance by various clinical isolates of *Candida* species is unknown.

The prevalence of resistance in *Aspergillus* spp and the exact mechanism are not known.

NONCLINICAL TOXICOLOGY

Carcinogenesis, Mutagenesis, Impairment Of Fertility

Hepatic carcinomas and adenomas were observed in a 6-month intravenous toxicology study with an 18-month recovery period of micafungin sodium in rats designed to assess the reversibility of hepatocellular lesions.

Rats administered micafungin sodium for 3 months at 32 mg/kg/day (corresponding to 8 times the highest recommended human dose [150 mg/day], based on AUC comparisons), exhibited colored patches/zones, multinucleated hepatocytes and altered hepatocellular foci after 1 or 3 month recovery periods, and adenomas were observed after a 21-month recovery period. Rats administered micafungin sodium at the same dose for 6 months exhibited adenomas after a 12-month recovery period; after an 18-month recovery period, an increased incidence of adenomas was observed, and additionally, carcinomas were detected. A lower dose of micafungin sodium (equivalent to 5 times the human AUC) in the 6-month rat study resulted in a lower incidence of adenomas and carcinomas following 18 months recovery. The duration of micafungin dosing in these rat studies (3 or 6 months) exceeds the usual duration of *MYCAMINE* dosing in patients, which is typically less than 1 month for treatment of esophageal candidiasis, but dosing may exceed 1 month for *Candida* prophylaxis.

Although the increase in carcinomas in the 6-month rat study did not reach statistical significance, the persistence of altered hepatocellular foci subsequent to micafungin dosing, and the presence of adenomas and carcinomas in the recovery periods suggest a causal relationship between micafungin sodium, altered hepatocellular foci, and hepatic neoplasms. Whole-life carcinogenicity studies of *MYCAMINE* in animals have not been conducted, and it is not known whether the hepatic neoplasms observed in treated rats also occur in other species, or if there is a dose threshold for this effect.

Micafungin sodium was not mutagenic or clastogenic when evaluated in a standard battery of *in-vitro* and *in-vivo* tests (i.e., bacterial reversion - *S. typhimurium*, *E. coli*; chromosomal aberration; intravenous mouse micronucleus).

Male rats treated intravenously with micafungin sodium for 9 weeks showed vacuolation of the epididymal ductal epithelial cells at or above 10 mg/kg (about 0.6 times the recommended clinical dose for esophageal candidiasis, based on body surface area comparisons). Higher doses (about twice the recommended clinical dose, based on body surface area comparisons) resulted in higher epididymis weights and reduced numbers of sperm cells. In a 39-week intravenous study in dogs, seminiferous tubular atrophy and decreased sperm in the epididymis were observed at 10 and 32 mg/kg, doses equal to about 2 and 7 times the recommended clinical dose, based on body surface area comparisons. There was no impairment of fertility in animal studies with micafungin sodium.

Animal Toxicology And/Or Pharmacology

High doses of micafungin sodium (5 to 8 times the highest recommended human dose, based on AUC comparisons) have been associated with irreversible changes to the liver when administered for 3 or 6 months, and these changes may be indicative of pre-malignant processes (see *Nonclinical Toxicology*).

Reproductive Toxicology Studies

Micafungin sodium administration to pregnant rabbits (intravenous dosing on days 6 to 18 of gestation) resulted in visceral abnormalities and abortion at 32 mg/kg, a dose equivalent to about four times the recommended dose based on body surface area comparisons. Visceral abnormalities included abnormal lobation of the lung, levocardia, retrocaval ureter, anomalous right subclavian artery, and dilatation of the ureter.

CLINICAL STUDIES

Treatment of Candidemia and Other Candida Infections

Two dose levels of *MYCAMINE* were evaluated in a randomized, double-blind study to determine the efficacy and safety versus caspofungin in patients with invasive candidiasis and candidemia. Patients were randomized to receive once daily intravenous infusions (IV) of *MYCAMINE*, either 100 mg/day or 150 mg/day or caspofungin (70 mg loading dose followed by 50 mg maintenance dose). Patients in both study arms were permitted to switch to oral fluconazole after at least 10 days of intravenous therapy, provided they were non-neutropenic, had improvement or resolution of clinical signs and symptoms, had a *Candida* isolate which was susceptible to fluconazole, and had documentation of 2 negative cultures drawn at least 24 hours apart. Patients were stratified by APACHE II score (≤ 20 or >20) and by geographic region. Patients with *Candida* endocarditis were excluded from this analysis. Outcome was assessed by overall treatment success based on clinical (complete resolution or improvement in attributable signs and symptoms and radiographic abnormalities of the *Candida* infection and no additional antifungal therapy) and mycological (eradication or presumed eradication) response at the end of IV therapy. Deaths that occurred during IV study drug therapy were treated as failures.

In this study, 111/578 (19.2 %) of the patients had baseline APACHE II scores of >20 , and 50/578 (8.7%) were neutropenic at baseline (absolute neutrophil count less than 500 cells/mm³). Outcome, relapse and mortality data are shown for the recommended dose of *MYCAMINE* (100 mg/day) and caspofungin in Table 7.

Table 7: Efficacy Analysis: Treatment Success in Patients in Study 03-0-192 with Candidemia and other *Candida* Infections

	<i>MYCAMINE</i> 100 mg/day n (%) % treatment difference (95%CI)	Caspofungin 70/50 mg/day¹ n (%)
Treatment Success at End of IV Therapy²	135/191 (70.7) 7.4 (-2.0, 16.3)	119/188 (63.3)
Success in Patients with Neutropenia at Baseline	14/22 (63.6)	5/11 (45.5)
Success by Site of Infection		
Candidemia	116/163 (71.2)	103/161 (64)
Abscess	4/5 (80)	5/9 (55.6)
Acute Disseminated³	6/13 (46.2)	5/9 (55.6)
Endophthalmitis	1/3	1/1
Chorioretinitis	0/3	0
Skin	1/1	0
Kidney	2/2	1/1
Pancreas	1/1	0
Peritoneum	1/1	0
Lung/Skin	0/1	0
Lung/Spleen	0/1	0
Liver	0	0/2
Intraabdominal abscess	0	3/5

Chronic Disseminated Peritonitis	0/1 4/6 (66.7)	0 2/5 (40)
Success by Organism ⁴		
<i>C. albicans</i>	57/81 (70.4)	45/73 (61.6)
<i>C. glabrata</i>	16/23 (69.6)	19/31 (61.3)
<i>C. tropicalis</i>	17/27 (63)	22/29 (75.9)
<i>C. parapsilosis</i>	21/28 (75)	22/39 (56.4)
<i>C. krusei</i>	5/8 (62.5)	2/3 (66.7)
<i>C. guilliermondii</i>	1/2	0/1
<i>C. lusitaniae</i>	2/3 (66.7)	2/2
Relapse through 6 Weeks ⁵		
Overall	49/135 (36.3)	44/119 (37)
Culture confirmed relapse	5	4
Required systemic antifungal therapy	11	5
Died during follow-up	17	16
Not assessed	16	19
Overall study mortality	58/200 (29)	51/193 (26.4)
Mortality during IV therapy	28/200 (14)	27/193 (14)

¹70 mg loading dose on day 1 followed by 50 mg/day thereafter (caspofungin)

²All patients who received at least one dose of study medication and had documented invasive candidiasis or candidemia. Patients with *Candida* endocarditis were excluded from the analyses.

³A patient may have had >1 organ of dissemination.

⁴A patient may have had >1 baseline infection species.

⁵All patients who had a culture confirmed relapse or required systemic antifungal therapy in the post treatment period for a suspected or proven *Candida* infection. Also includes patients who died or were not assessed in follow-up.

In two cases of ophthalmic involvement assessed as failures in the above table due to missing evaluation at the end of IV treatment with *MYCAMINE*, therapeutic success was documented during protocol-defined oral fluconazole therapy.

Treatment of Esophageal Candidiasis

In two controlled trials involving 763 patients with esophageal candidiasis, 445 adults with endoscopically-proven candidiasis received *MYCAMINE*, and 318 received fluconazole for a median duration of 14 days (range 1-33 days).

MYCAMINE was evaluated in a randomized, double-blind study which compared *MYCAMINE* 150 mg/day (n=260) to intravenous fluconazole 200 mg/day (n=258) in adults with endoscopically-proven esophageal candidiasis. Most patients in this study had HIV infection, with CD4 cell counts <100 cells/mm³. Outcome was assessed by endoscopy and by clinical response at the end of treatment. Endoscopic cure was defined as endoscopic grade 0, based on a scale of 0-3. Clinical cure was defined as complete resolution in clinical symptoms of esophageal candidiasis (dysphagia, odynophagia, and retrosternal pain). Overall therapeutic cure was defined as both clinical and endoscopic cure. Mycological eradication was determined by culture, and by histological or cytological evaluation of esophageal biopsy or brushings obtained endoscopically at the end of treatment. As shown in Table 8,

endoscopic cure, clinical cure, overall therapeutic cure, and mycological eradication were comparable for patients in the *MYCAMINE* and fluconazole treatment groups.

Table 8: Endoscopic, Clinical, and Mycological Outcomes for Esophageal Candidiasis at End-of-Treatment

Treatment Outcome*	<i>MYCAMINE</i> 150 mg/day n=260	Fluconazole 200 mg/day n=258	% Difference† (95% CI)
Endoscopic Cure	228 (87.7%)	227 (88.0%)	-0.3% (-5.9, +5.3)
Clinical Cure	239 (91.9%)	237 (91.9%)	0.06% (-4.6, +4.8)
Overall Therapeutic Cure	223 (85.8%)	220 (85.3%)	0.5% (-5.6, +6.6)
Mycological Eradication	141/189 (74.6%)	149/192 (77.6%)	-3.0% (-11.6, +5.6)

*Endoscopic and clinical outcome were measured in modified intent-to-treat population, including all randomized patients who received ≥ 1 dose of study treatment. Mycological outcome was determined in the per protocol (evaluable) population, including patients with confirmed esophageal candidiasis who received at least 10 doses of study drug, and had no major protocol violations.

† Calculated as *MYCAMINE* – fluconazole

Most patients (96%) in this study had *Candida albicans* isolated at baseline. The efficacy of *MYCAMINE* was evaluated in less than 10 patients with *Candida* species other than *C. albicans*, most of which were isolated concurrently with *C. albicans*.

Relapse was assessed at 2 and 4 weeks post-treatment in patients with overall therapeutic cure at end of treatment. Relapse was defined as a recurrence of clinical symptoms or endoscopic lesions (endoscopic grade > 0). There was no statistically significant difference in relapse rates at either 2 weeks or through 4 weeks post-treatment for patients in the *MYCAMINE* and fluconazole treatment groups, as shown in Table 9.

Table 9: Relapse of Esophageal Candidiasis at Week 2 and through Week 4 Post-Treatment in Patients with Overall Therapeutic Cure at the End of Treatment

Relapse	<i>MYCAMINE</i> 150 mg/day n=223	Fluconazole 200 mg/day n=220	% Difference* (95% CI)
Relapse † at Week 2	40 (17.9%)	30 (13.6%)	4.3% (-2.5, 11.1)
Relapse† Through Week 4 (cumulative)	73 (32.7%)	62 (28.2%)	4.6% (-4.0, 13.1)

*Calculated as *MYCAMINE* – fluconazole; N=number of patients with overall therapeutic cure (both clinical and endoscopic cure at end-of-treatment)

† Relapse included patients who died or were lost to follow-up, and those who received systemic anti-fungal therapy in the post-treatment period

In this study, 459 of 518 (88.6%) patients had oropharyngeal candidiasis in addition to esophageal candidiasis at baseline. At the end of treatment 192/230 (83.5%) *MYCAMINE* treated patients and 188/229 (82.1%) of fluconazole treated patients experienced resolution of

signs and symptoms of oropharyngeal candidiasis. Of these, 32.3% in the *MYCAMINE* group, and 18.1% in the fluconazole group (treatment difference = 14.2%; 95% confidence interval [5.6, 22.8]) had symptomatic relapse at 2 weeks post-treatment. Relapse included patients who died or were lost to follow-up, and those who received systemic antifungal therapy during the post-treatment period. Cumulative relapse at 4 weeks post-treatment was 52.1% in the *MYCAMINE* group and 39.4% in the fluconazole group (treatment difference 12.7%, 95% confidence interval [2.8, 22.7]).

Prophylaxis of Candida or Aspergillus Infections in Hematopoietic Stem Cell Transplant Recipients

In a randomized, double-blind study, *MYCAMINE* (50 mg IV once daily) was compared to fluconazole (400 mg IV once daily) in 882 patients undergoing an autologous or syngeneic (46%) or allogeneic (54%) stem cell transplant. The status of the patients' underlying malignancy at the time of randomization was: 365 (41%) patients with active disease, 326 (37%) patients in remission, and 195 (22%) patients in relapse. The more common baseline underlying diseases in the 476 allogeneic transplant recipients were: chronic myelogenous leukemia (22%), acute myelogenous leukemia (21%), acute lymphocytic leukemia (13%), and non-Hodgkin's lymphoma (13%). In the 404 autologous and syngeneic transplant recipients the more common baseline underlying diseases were: multiple myeloma (37.1%), non-Hodgkin's lymphoma (36.4%), and Hodgkin's disease (15.6%). During the study, 198 of 882 (22.4%) transplant recipients had proven graft-versus-host disease; and 475 of 882 (53.9%) recipients received immunosuppressive medications for treatment or prophylaxis of graft-versus-host disease.

Study drug was continued until the patient had neutrophil recovery to an absolute neutrophil count (ANC) of ≥ 500 cells/mm³ or up to a maximum of 42 days after transplant. The average duration of drug administration was 18 days (range 1 to 51 days).

Successful prophylaxis was defined as the absence of a proven, probable, or suspected systemic fungal infection through the end of therapy (usually 18 days), and the absence of a proven or probable systemic fungal infection through the end of the 4-week post-therapy period. A suspected systemic fungal infection was diagnosed in patients with neutropenia (ANC < 500 cells/mm³); persistent or recurrent fever (while ANC < 500 cells/mm³) of no known etiology; and failure to respond to at least 96 hours of broad spectrum antibacterial therapy. A persistent fever was defined as four consecutive days of fever greater than 38°C. A recurrent fever was defined as having at least one day with temperatures ≥ 38.5 °C after having at least one prior temperature > 38 °C; or having two days of temperatures > 38 °C after having at least one prior temperature > 38 °C. Transplant recipients who died or were lost to follow-up during the study were considered failures of prophylactic therapy.

Successful prophylaxis was documented in 80.7% of recipients who received *MYCAMINE*, and in 73.7% of recipients who received fluconazole (7.0% difference [95% CI = 1.5, 12.5]), as shown in Table 10, along with other study endpoints. The use of systemic antifungal therapy post-treatment was 42% in both groups.

The number of proven breakthrough *Candida* infections was 4 in the *MYCAMINE* and 2 in the fluconazole group.

The efficacy of *MYCAMINE* against infections caused by fungi other than *Candida* sp. and *Aspergillus* sp. has not been established.

Table 10: Results from Clinical Study of Prophylaxis of Candida Infections in Hematopoietic Stem Cell Transplant Recipients

Outcome of Prophylaxis	<i>MYCAMINE</i> 50 mg/day (n=425)	Fluconazole 400 mg/day (n=457)
Success *	343 (80.7%)	337 (73.7%)
Failure:	82 (19.3%)	120 (26.3%)
All Deaths ¹	18 (4.2%)	26 (5.7%)
Proven/probable fungal infection prior to death	1 (0.2%)	3 (0.7%)
Proven/probable fungal infection (not resulting in death) ¹	6 (1.4%)	8 (1.8%)
Suspected fungal infection ²	53 (12.5%)	83 (18.2%)
Lost to follow-up	5 (1.2%)	3 (0.7%)

*Difference (*MYCAMINE* – Fluconazole): +7.0% [95% CI=1.5, 12.5]

¹Through end-of-study (4 weeks post- therapy)

²Through end-of-therapy

Treatment of Aspergillosis

The therapeutic efficacy and safety of micafungin for the treatment of aspergillosis have been evaluated in a number of studies. The effectiveness of micafungin monotherapy in different types of *Aspergillus* infections are mentioned in the tables below.

Table 11: Result of a clinical study conducted in Japan

Strain	Disease	Duration of treatment (days) Median (min.-max.)	Effective cases/total	Effectiveness (%)
<i>Aspergillus</i>	Invasive pulmonary aspergillosis	26 (8-56)	6/10	60.0
	Chronic necrotizing pulmonary aspergillosis	56 (11-57)	6/8	-
	Pulmonary aspergilloma	39.5 (18-56)	10/16	62.5
	Sub total	39.5 (8-57)	22/34	64.7

Table 12: Result of a clinical study conducted in countries other than Japan

Strain	Disease	Duration of treatment (days) Median (min.-max.)	Effective cases/total	Effectiveness (%)
<i>Aspergillus</i>	Invasive pulmonary aspergillosis	34 (7-126)	32/39	82.1
	Paranasal sinus aspergillosis (suspected)	17 (14-29)	1/3	-
	Aspergillemia	17	1/1	-
	Sub total	33 (7-126)	34/43	79.1

PHARMACEUTICAL PARTICULARS**List of Excipients**

Lactose monohydrate
Citric acid anhydrous (to adjust the pH)
Sodium hydroxide (to adjust the pH)

Incompatibilities

This medicinal product must not be mixed or co-infused with other medicinal products except those mentioned in *Posology and Method of Administration (Directions for Reconstitution and Dilution)*.

Shelf Life

Unopened vial: 3 years.

The expiry date is indicated on the label and packaging.

Reconstituted concentrate in vial:

Chemical and physical in-use stability has been demonstrated for up to 48 hours at 25°C when reconstituted with 0.9% Sodium Chloride solution for infusion or 5% Dextrose solution for infusion.

Diluted infusion solution:

Chemical and physical in-use stability has been demonstrated for 96 hours at 25°C when protected from light when diluted with 0.9% Sodium Chloride solution for infusion or 5% Dextrose solution for infusion.

MYCAMINE contains no preservatives.

From a microbiological point of view, the reconstituted and diluted solutions should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the

responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless the reconstitution and dilution have taken place in controlled and validated aseptic conditions.

Special Precautions for Storage

Unopened vials: Store at or below 25°C. Protect from light.

For storage conditions of the reconstituted and diluted medicinal product, see *Shelf Life*.

Keep out of reach of children.

Nature and Specification of Container

10 ml Type I glass vial with an isobutylene-isoprene (Teflon-laminated) rubber stopper and a flip-off cap. The vial is wrapped with an UV-protective film.

Supplied in packs of 1 vial

All presentations may not be marketed in India.

Instructions for Use / Handling

- Any unused product or waste material should be disposed of in accordance with local requirements.
- *MYCAMINE* must not be mixed or co-infused with other medicinal products except those mentioned under section *Posology and Method of Administration (Directions for Reconstitution and Dilution)*.
- The vial is for single use only. Therefore, please discard unused reconstituted concentrate immediately.

Manufactured by:

Astellas Pharma Tech Co., Ltd.
Takaoka Plant
30 Toidesakae-machi,
Takaoka City, Toyama 939-1118,
Japan

Imported by:

Astellas Pharma India Pvt. Ltd.,
301, 3rd Floor, C and B Square,
127, Andheri-Kurla Road, Chakala,
Andheri (East) Mumbai
Tal: Andheri East (Mumbai – Zone 6) 400069

Marketed by:

GlaxoSmithKline Pharmaceuticals Limited.
(Under licence from Astellas Pharma Inc., Japan)
Regd. Office:

Dr. Annie Besant Road, Worli, Mumbai 400030, India.

® Registered Trade Mark of Astellas Pharma Inc. Japan

Version: MYC/PI/IN/2013/01 dated 10-Jul-13

*Adapted from USPI dated Jun 2011 and certain sections of EU SPC dated 20-Dec-2010
(Ref: Astellas Pharma Inc., Micafungin CCDS version 1.6 dated 07-Mar-2011]*